


D5



	Very Poorly Drained
	Poorly Drained
	Poorly and Very Poorly Drained
	Alluvial and Floodplain

-  Parcels
 Rivers/Streams
 Reservoir/Lake/Pond
 Swamp

55 Map Unit (Soil Series)

I. Glacial Till Soils - Soils formed in unstratified sand, silt and rock derived from materials weathered from schist, granite and gneiss

- 2 Ridgebury fine sandy loam
- 3 Ridgebury, Leicester, and Whitman soils, extremely stony
- 4 Leicester fine sandy loam

II. Glaciofluvial Soils - Soils formed in stratified sand and gravel derived from acid crystalline rocks and deposited by glacial meltwater

- | | |
|----|--------------------|
| 12 | Raypol silt loam |
| 13 | Walpole sandy loam |
| 15 | Scarboro muck |

III. Flooplain Soils - Soils formed in alluvium deposited by streams and


- | | |
|-----|---|
| 100 | Suncook loamy fine sand |
| 101 | Occum fine sandy loam |
| 102 | Pootatuck fine sandy loam |
| 103 | Rippowam fine sandy loam |
| 108 | Saco silt loam |
| 109 | Fluvaquents-Udifluvents complex, frequently flooded |
| 301 | Beaches-Udipsamments complex, coastal |

IV. **Organic Soils** - Soils formed in shallow to deep organic deposits

- 17 Timakwa and Natchaug soils
18 Catden and Freetown soils

V. **Disturbed Wetland Soils** - Soils which have had two or more feet of their original soil surface disturbed through fillinf, excavation or grading; have a wetlnad hydrologi regime; and are capable of supporting wetland vegetation

- ## 309 Aquents



1 inch = 200 feet

MAP SCALE & ACCURACY:
Soils mapped at a scale of 1:12,000 (1 inch = 1,000 feet)
Minimum soil map unit delineation is 3 acres in size

MAP REFERENCES
Planimetrics Data Source: AT&T/SBC
Parcel Data Source: Town of East Hampton
Soils Data Source: USDA, NRCS Soil Survey Geographic (SSURGO) database for the State of Connecticut.

F3	F4	F5
E3	E4	E5
D3	D4	D5